

**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings of claims in the application:

**Listing of Claims:**

1-7. (Canceled)

1                   8.       (Currently amended) A method of visibly reducing a human skin wrinkle  
2 comprising: topically applying to the human skin wrinkle an imiquimod composition-IRM  
3 ~~compound that is an agonist of TLR7, TLR8, or both TLR7 and TLR8~~ in an amount and for a  
4 period of time sufficient to visibly reduce the wrinkle, wherein imiquimod is the sole active  
5 pharmaceutical ingredient applied to the human skin wrinkle in performance of the method;  
6 ~~wherein the IRM compound is an imidazoquinoline amine, a tetrahydroimidazoquinoline amine,~~  
7 ~~an imidazopyridine amine, a 1,2-bridged imidazoquinoline amine, a 6,7-fused~~  
8 ~~cycloalkylimidazopyridine amine, an imidazonaphthyridine amine, a~~  
9 ~~tetrahydroimidazonaphthyridine amine, an oxazoloquinoline amine, a thiazoloquinoline amine,~~  
10 ~~an oxazolopyridine amine, a thiazolopyridine amine, an oxazonaphthyridine amine, a~~  
11 ~~thiazolonaphthyridine amine, or a combination thereof.~~

9-32. (Canceled)

1                   33.       (New) The method of claim 8, wherein the imiquimod composition is  
2 applied daily.

1                   34.       (New) The method of claim 8, wherein the imiquimod composition is  
2 applied once per day two to three times a week.

1                   35.       (New) The method of claim 8, wherein the imiquimod composition  
2 comprises 5% imiquimod by weight.

1                   36.     (New) The method of claim 8, wherein the imiquimod composition about  
2     1% to about 2.5% by weight.

1                   37.     (New) The method of claim 8, wherein the imiquimod composition is  
2     applied to the human skin wrinkle for about one month.

1                   38.     (New) The method of claim 8, wherein the imiquimod composition is  
2     applied to the human skin wrinkle for about two months.

1                   39.     (New) The method of claim 8, wherein the imiquimod composition is  
2     applied to the human skin wrinkle for about 4 to 12 months.

1                   40.     (New) The method of claim 8, further comprising measuring the  
2     reduction in human skin wrinkles by visual or photographic examination of the skin.

1                   41.     (New) The method of claim 8, wherein the imiquimod composition is  
2     about 5% imiquimod by weight, and the imiquimod composition is applied to the human skin  
3     wrinkle two to three times a week.

1                   42.     (New) The method of claim 8, wherein the imiquimod composition is  
2     about 1% imiquimod by weight, and the imiquimod composition is applied daily to the human  
3     skin wrinkle.

1                   43.     (New) The method of claim 8, wherein the imiquimod composition is  
2     about 2.5% imiquimod by weight, and the imiquimod composition is applied daily to the human  
3     skin wrinkle.